



Measure 1: Diabetes: Hemoglobin A1c Poor Control

Reporting Option: Claims, Diabetic Retinopathy Measures Group via Registry (Cross-cutting Measure)

Quality Domain: Effective Clinical Care

Instructions: This measure is to be reported a minimum of once per reporting period for patients with diabetes seen during the reporting period. The most recent quality-data code submitted will be used for performance calculation. This measure may be reported by clinicians who perform the quality actions described in the measure based on the services provided and the measure-specific denominator coding.

Category II Codes:

Numerator Instructions: A lower calculated performance rate for this measure indicates better clinical care or control. Patient is numerator compliant if most recent HbA1c level >9% or is missing a result or if an HbA1c test was not done during the measurement year.

3064F Most recent hemoglobin A1c level > 9.0%

Or

3046F 8P Hemoglobin A1c level was not performed during the performance period (12 months)

Or

3044F Most recent hemoglobin A1c (HbA1c) level < 7.0%

Or

3045F Most recent hemoglobin A1c (HbA1c) level 7.0 to 9.0%

CPT codes:

97802, 97803, 97804, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99217, 99218, 99219, 99220, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99281, 99282, 99283, 99284, 99285, 99291, 99304, 99305, 99306, 99307, 99308, 99309, 99310, 99315, 99316, 99318, 99324, 99325, 99326, 99327, 99328, 99334, 99335, 99336, 99337, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, G0270, G0271, G0402, G0438, G0439

Diagnosis Codes:

E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.321, E10.329, E10.331, E10.339, E10.341, E10.349, E10.351, E10.359, E10.36, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9, E11.00, E11.01, E11.21, E11.22, E11.29, E11.311, E11.319, E11.321, E11.329, E11.331,



E11.339, E11.341, E11.349, E11.351, E11.359, E11.36, E11.39, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9, E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.321, E13.329, E13.331, E13.339, E13.341, E13.349, E13.351, E13.359, E13.36, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9, O24.011, O24.012, O24.013, O24.019, O24.02, O24.03, O24.111, O24.112, O24.113, O24.119, O24.12, O24.13, O24.311, O24.312, O24.313, O24.319, O24.32, O24.33, O24.811, O24.812, O24.813, O24.819, O24.82, O24.83

Rationale: Diabetes mellitus (diabetes) is a group of diseases characterized by high blood glucose levels caused by the body's inability to correctly produce or utilize the hormone insulin. It is recognized as a leading cause of death and disability in the U.S. and is highly underreported as a cause of death. Diabetes may cause life-threatening, life ending or life-altering complications, including poor circulation, nerve damage or neuropathy in the feet and eventual amputation. Nearly 60-70 percent of diabetics suffer from mild or severe nervous system damage (American Diabetes Association 2009).

Randomized clinical trials have demonstrated that improved glycemic control, as evidenced by reduced levels of glycohemoglobin, correlates with a reduction in the development of microvascular complications in both Type 1 and Type 2 diabetes (Diabetes Control and Complications Trial Research Group 1993; Ohkubo 1995). In particular, the Diabetes Control and Complications Trial (DCCT) showed that for patients with Type 1 diabetes mellitus, important clinical outcomes such as retinopathy (an important precursor to blindness), nephropathy (which precedes renal failure), and neuropathy (a significant cause of foot ulcers and amputation in patients with diabetes) are directly related to level of glycemic control (Diabetes Control and Complications Trial Research Group 1993). Similar reductions in complications were noted in a smaller study of intensive therapy of patients with Type 2 diabetes by Ohkubo and co-workers, which was conducted in the Japanese population (Ohkubo et al. 1995).

Clinical Recommendation Statements: For frail older adults, persons with life expectancy of less than 5 years, and others in whom the risks of intensive glycemic control appear to outweigh the benefits, a less stringent target such as 8% is appropriate. (Quality of Evidence: Level III; Strength of Evidence: Grade B)