

QCDR Measure:

IRIS-18: Chronic Anterior Uveitis: Post-treatment visual acuity

National Quality Strategy Domain:

Effective Clinical Care

Measure Type:

Outcome

Description:

Percentage of chronic anterior uveitis patients with a post-treatment best corrected visual acuity of 20/40 or greater OR patients whose visual acuity had returned to their baseline value prior to onset of uveitis.

Instructions:

This measure is to be reported a minimum of once per reporting period for patients seen during the reporting period. It is anticipated that clinicians who provide the primary management of patients with chronic anterior uveitis (in either one or both eyes) will submit this measure.

Denominator:

All patients aged 18 years or greater who underwent treatment for chronic anterior uveitis.

Patients aged ≥ 18 years

AND

Diagnosis of Chronic Anterior Uveitis

ICD-9 [for use 1/1/2015 – 9/30/2015]

- Chronic iridocyclitis, unspecified (ICD-9: 364.10)
- Chronic iridocyclitis in diseases classified elsewhere (ICD-9: 364.11)

ICD-10 [for use 10/1/2015 – 12/31/2015]

- Chronic iridocyclitis (ICD-10: H20.10, H20.11, H20.12, H20.13)

AND

Treatment for Acute Uveitis which includes topical corticosteroids and antimicrobial/antivirals for infectious cases.

Numerator:

Patients with a best corrected visual acuity of 20/40 or greater within 90 days of treatment initiation

OR

Patients whose visual acuity had returned to their baseline value prior to onset of acute uveitis within 90 days of treatment initiation

Numerator Options:

Performance Met: Best Corrected visual acuity of 20/40 or better achieved within 90 days following treatment initiation

OR

Patient's visual acuity returned to baseline value within 90 days of treatment initiation

Performance Not Met: Patient did not return to baseline OR achieve a best corrected visual acuity of 20/40 within 90 days following treatment initiation

Improvement Notation:

Higher score indicates better performance