Effective Clinical Care
Measure Type:
Outcome
Description:
Percentage of patients with nonexudative age-related macular degeneration progressing to exudative age-related macular degeneration over the past 12 months.
Instructions:
This measure is to be reported a minimum of <u>once per reporting period</u> for patients seen during the reporting period. It is anticipated that clinicians who provide the primary management of patients with exudative age-related macular degeneration (in either one or both eyes) will submit this measure.
Denominator:
All patients aged 18 years of older with an active diagnosis of exudative age-related macular degeneration including documentation of the laterality (OD, OS, OU)
Denominator Criteria
Patients aged ≥ 18 years
AND
Diagnosis of Exudative Age-Related Macular Degeneration
ICD-9 [for use 1/1/2015 – 9/30/2015]
• Exudative age-related macular degeneration (ICD-9: 362.52)

Numerator:

QCDR Measure:

National Quality Strategy Domain:

IRIS-12: Age-Related Macular Degeneration: Disease Progression

Patients who had an active diagnosis of nonexudative age-related macular degeneration in the past 12 months including documentation of the laterality (OD, OS, OU) and a history of taking AREDS supplements

Exudative age-related macular degeneration (ICD-10: H35.32)

Revised 04/20/2015

ICD-10 [for use 10/1/2015 – 12/31/2015]

Numerator Options:

Performance Met: Patients who advanced from nonexudative AMD to exudative

AMD in the past 12 months

Performance Not Met: Patients who did not advanced from nonexudative AMD to

exudative AMD in the past 12 months

Patients are only eligible for the numerator when the disease advancement occurs in the same eye.

Improvement Notation:

Lower score indicates better performance