



Measure 130: Documentation of Current Medications in the Medical Record

Reporting Options: Claims, Registry, Cataracts Measures Group, Diabetic Retinopathy Measures Group (Cross-cutting Measure)

Quality Domain: Patient Safety

Description: Percentage of patients aged 18 years and older with a list of all current medications (includes prescription, over-the-counter, herbals, vitamin/mineral/dietary nutritional supplements) documented by the provider, including drug name, dosage, frequency and route.

Instructions: This measure is to be reported each visit during the 12 month reporting period. Eligible professionals meet the intent of this measure by making their best effort to document a current, complete and accurate medication list during each encounter. There is no diagnosis associated with this measure. This measure may be reported by eligible professionals who perform the quality actions described in the measure based on the services provided and the measure-specific denominator coding.

Category II Codes:

G8427 List of current medications with name, dosages, frequency and route documented;

Or

G8430 Provider documentation that patient is not eligible for medication assessment;

Or

G8428 Current medications with name, dosages, frequency, route not documented, reason not specified

CPT Codes:

90791, 90792, 90832, 90834, 90837, 90839, 90957, 90958, 90959, 90960, 90962, 90965, 90966, 92002, 92004, 92012, 92014, 92507, 92508, 92526, 92541, 92542, 92543, 92544, 92545, 92547, 92548, 92557, 92567, 92568, 92570, 92585, 92588, 92626, 96116, 96150, 96151, 96152, 97001, 97002, 97003, 97004, 97532, 97802, 97803, 97804, 98960, 98961, 98962, 99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99221, 99222, 99223, 99324, 99325, 99326, 99327, 99328, 99334, 99335, 99336, 99337, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99495, 99496, G0101, G0108, G0270, G0402, G0438, G0439

Diagnosis Codes



None required for this measure. If your computer system requires a diagnosis, it is appropriate to append the associated exam diagnosis.

Clinical Recommendation Statements:

The Joint Commission's 2015 Ambulatory Care National Patient Safety Goals guide providers to maintain and communicate accurate patient medication information. Specifically, the section "Use Medicines Safely NPSG.03.06.01" states the following: "Maintain and communicate accurate patient medication information. The types of information that clinicians use to reconcile medications include (among others) medication name, dose, frequency, route, and purpose. Organizations should identify the information that needs to be collected to reconcile current and newly ordered medications and to safely prescribe medications in the future." (Joint Commission, 2015, retrieved at: National Patient Safety Goals Effective January 1, 2015).

The National Quality Forum's 2010 update of the Safe Practices for Better Healthcare, states healthcare organizations must develop, reconcile, and communicate an accurate patient medication list throughout the continuum of care. Improving the safety of healthcare delivery saves lives, helps avoid unnecessary complications, and increases the confidence that receiving medical care actually makes patients better, not worse. Every healthcare stakeholder group should insist that provider organizations demonstrate their commitment to reducing healthcare error and improving safety by putting into place evidence-based safe practices.

The AMA's published report, *The Physician's Role in Medication Reconciliation*, identified the best practice medication reconciliation team as one that is multidisciplinary and—in all settings of care—will include physicians, pharmacists, nurses, ancillary health care professionals and clerical staff. The team's variable requisite knowledge, skills, experiences, and perspectives are needed to make medication reconciliation work as safely and smoothly as possible. Team members may have access to vital information or data needed to optimize medication safety. Because physicians are ultimately responsible for the medication reconciliation process and subsequently accountable for medication management, physician leadership and involvement in all phases of developing and initiating a medication reconciliation process or model is important to its success.