

**QCDR Measure:**

AAO-5: Surgery for Acquired Involutional Ptosis: Patients with an improvement of marginal reflex distance (MRD)

National Quality Strategy Domain:

Effective Clinical Care

Measure Type:

Outcome

Description:

Percentage of surgical ptosis patients with an improvement of MRD postoperatively.

Instructions:

This measure is to be reported **each time** a ptosis surgical procedure is performed during the reporting period. The measure is intended to reflect the quality of services provided by the surgeon performing the procedure.

Denominator:

All patients aged 18 years or older with a diagnosis of acquired involutional ptosis who underwent a surgical procedure for the condition.

Denominator Criteria

Patients aged ≥ 18 years

AND

Diagnosis of Involutional Ptosis

Myogenic Ptosis (ICD-10: H02.421, H02.422, H02.423, H02.429)

AND

Ptosis Surgical Procedure

- Internal levator resection (CPT: 67903)
- External levator resection (CPT: 67904)
- Conjunctivo-tarso-Muller's (CPT: 67908)

Numerator:

Patients with an improvement of MRD postoperatively compared to preoperative level.

Numerator Options:

Performance Met:	Patients who achieved an improvement in MRD postoperatively within 90 days following surgery
Performance Not Met:	Patients who did not achieve an improvement in MRD postoperatively within 90 days following surgery

Improvement Notation:

Higher score indicates better performance