

## **POLICY STATEMENT**

### **Laser Surgery**

#### **Policy:**

The quality of care of patients undergoing laser surgery is a concern of the American Academy of Ophthalmology. This quality of care should be safeguarded in the same way that it is for patients undergoing any surgery because the clinical complexities and potentially harmful consequences are similar. The Academy strongly supports the position that all laser surgery for medical purposes, including ophthalmic laser surgery, should be performed only by licensed doctors of medicine or osteopathy.

#### **Background:**

Like other surgical procedures, laser surgery alters, removes, replaces, and reshapes human tissue or it activates drugs for the purpose of treating disease and improving patient function and well being. Because laser energy possesses remarkable strength, unlike ordinary light energy, lasers are used as sophisticated surgical instruments to produce definitive and precise surgical effects.

Lasers were originally used in ophthalmology in the early 1960s soon after the laser effect was first successfully demonstrated. Since then, many benefits of laser surgery have been demonstrated, including reduced visual loss from diabetic retinopathy and age-related macular degeneration, restored vision for posterior capsular opacification that occurs after cataract surgery, and reduced intraocular pressure in patients with glaucoma.

The excimer laser has been approved by the FDA for use in refractive surgery, under specific terms and conditions. Adverse effects of this surgery that have occurred include corneal haze, which can limit visual acuity, and, rarely, infection or scarring. The most recently FDA-approved ophthalmic laser procedure, photodynamic therapy, was introduced in ophthalmic practice in the spring of 2000. This application, which uses laser light to activate an intravenously administered drug, represents a significant advance for some patients with certain patterns of choroidal neovascularization.

Like other surgical instruments, lasers are potentially dangerous and can cause bleeding, edema, trauma and tissue damage. Specific risks associated with lasers that are used for ophthalmic surgery include penetration of a blood vessel with resultant bleeding, increase in intraocular pressure with resultant damage to the optic nerve, retinal injury, damage to the cornea, and cataract formation. Each of these complications could result in significant loss of vision. Because of the potentially harmful medical consequences of laser surgery, strict guidelines have been established for the use of lasers in patient care. The FDA regulates all medical instruments as prescription devices, including ophthalmic lasers.

In order to guard against potential proliferation of unsafe or inappropriate use of lasers in patient care, several surgical specialty societies have developed guidelines or policies that set forth criteria for the medical qualifications of surgeons practicing laser surgery. The guidelines also specify hospital privileges for laser surgery, residency training, and continuing medical education courses.

Thus, laser surgery is subject to the same high standards of care that govern all other medical practices.

**Evaluation:**

Ophthalmologists have been among the principal pioneers and innovators in the field of laser surgery. They learn and gain mastery of laser surgery techniques through residency and fellowship training, continuing educational courses, didactic courses, and preceptorship opportunities. The best possible outcome for a patient after laser surgery depends on the physician making an accurate diagnosis, consideration of the patient's health status and visual needs, choice of an appropriate treatment given the spectrum of alternatives, timing of treatment, and the use of prescription drugs postoperatively. As with any sophisticated surgery, patient outcome is also determined by the technical skill, dexterity and the coordination needed to control the laser system during treatment. Optimization of the outcome depends on the correct and timely recognition and management of both the anticipated and unforeseen complications.

**Recommendation:**

The clinical complexities of ocular laser surgery and the associated potential for laser-induced health problems clearly justify adherence by regulatory authorities to a position requiring that laser surgery be performed only by licensed doctors of medicine or osteopathy.

Approved by:	Board of Directors, June 1990
Revised and Approved by:	Board of Trustees, June 1993
Revised and Approved by:	Board of Trustees, October 1996
Revised and Approved by:	Board of Trustees, February 2001
Revised and Approved by:	Board of Trustees, March, 2006

©2006 American Academy of Ophthalmology®  
P.O. Box 7424 / San Francisco, CA 94102 / 415.561.8500