

**QCDR Measure:**

AAO-11: Nonexudative Age-Related Macular Degeneration: Loss of Visual Acuity

National Quality Strategy Domain:

Effective Clinical Care

Measure Type:

Outcome

Description:

Percentage of patients with a diagnosis of nonexudative age-related macular degeneration and taking AREDS supplements with of loss of less than 0.3 logMar of visual acuity within the past 12 months.

Instructions:

This measure is to be reported a minimum of once per reporting period for patients seen during the reporting period. It is anticipated that clinicians who provide the primary management of patients with nonexudative age-related macular degeneration (in either one or both eyes) will submit this measure.

Denominator:

All patients aged 18 years of older with a diagnosis of nonexudative AMD and documentation of laterality (OD, OS, OU)

Denominator Criteria

Patients aged ≥ 18 years

AND

Diagnosis of nonexudative age-related macular degeneration

- Nonexudative age-related macular degeneration (ICD-10: H35.31)

AND

Taking antioxidant supplements

Numerator:

Patients with two or more recorded visual acuity values within the past 12 months; loss of visual acuity less than 0.3 logMar. Visual acuity of the eye with nonexudative AMD should be reported.

Numerator Options:

Performance Met:	Patients who achieved a loss in visual acuity of ≤ 0.3 logMar
Performance Not Met:	Patients who did not achieve a loss in visual acuity of ≤ 0.3 logMar

Improvement Notation:

Higher score indicates better performance